

Complete Genome Sequence of a Porcine Epidemic Diarrhea Virus from a Novel Outbreak in Belgium, January 2015

Sebastiaan Theuns,^a Nádia Conceição-Neto,^b Isaura Christiaens,^a Mark Zeller,^b Lowiese M. B. Desmarets,^a Inge D. M. Roukaerts,^a Delphine D. Acar,^a Elisabeth Heylen,^b Jelle Matthijnsens,^b Hans J. Nauwynck^a

Ghent University, Faculty of Veterinary Medicine, Department of Virology, Parasitology and Immunology, Laboratory of Virology, Merelbeke, Belgium^a; KU Leuven–University of Leuven, Department of Microbiology and Immunology, Laboratory for Clinical and Epidemiological Virology, Rega Institute for Medical Research, Leuven, Belgium^b

S.T., N.C.-N., and I.C. contributed equally to this work.

J.M. and H.J.N. share senior authorship.

Porcine epidemic diarrhea virus (PEDV) is a member of the family *Coronaviridae* and can cause severe outbreaks of diarrhea in piglets from different age groups. Here, we report the complete genome sequence (28,028 nt) of a PEDV strain isolated during a novel outbreak in Belgium.

Received 14 April 2015 Accepted 17 April 2015 Published 21 May 2015

Citation Theuns S, Conceição-Neto N, Christiaens I, Zeller M, Desmarets LMB, Roukaerts IDM, Acar DD, Heylen E, Matthijnsens J, Nauwynck HJ. 2015. Complete genome sequence of a porcine epidemic diarrhea virus from a novel outbreak in Belgium, January 2015. *Genome Announc* 3(3):e00506-15. doi:10.1128/genomeA.00506-15.

Copyright © 2015 Theuns et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/4.0/).

Address correspondence to Sebastiaan Theuns, sebastiaan.theuns@ugent.be.

Porcine epidemic diarrheic virus (PEDV) caused outbreaks of diarrhea in Europe from the 1970s to the 1990s, but has since been reported only sporadically in the last few decades (1, 2). From 2010, severe outbreaks with mortality have been identified in Asia, and since the spring of 2013, the virus was detected for the first time in U.S. swine herds (3–9). Genetically, these American isolates were highly related (99.5% nucleotide similarity) to Asian PEDV strains (8, 10). In January 2014, a novel PEDV variant strain OH851 was isolated in the United States, which was associated with milder disease symptoms (11). Similar strains contained several deletions and insertions in their spike genes and were therefore called INDEL strains (6). In May 2014, an outbreak of diarrhea occurred in fattening pigs on a German farm, and the isolated strains were genetically almost identical to U.S. strain OH851 (12).

An outbreak of diarrhea in fattening pigs occurred on a Belgian pig farm at the end of January 2015, and the presence of PEDV RNA in the feces of these pigs was demonstrated using an in house RT-qPCR assay targeting the nucleocapsid gene. This was the first confirmed PEDV case in Belgium in decades. Therefore, the complete genome of this novel isolate, BEL/15V010/2015, was unraveled by next-generation sequencing, in order to assess its genetic relation to other PEDV isolates circulating around the globe. Viral particles were purified, RNA was extracted using the QIAamp Viral RNA mini kit (Qiagen), and whole transcriptome amplification was performed (WTA2 kit, Sigma Aldrich) to generate cDNA. The sequencing library was prepared using the Nextera XT DNA library preparation kit (Illumina). Sequencing was performed on a HiSeq 2500 platform (Illumina) for 301 cycles (150-bp paired-end reads). Raw reads were trimmed for quality and adapters using Trimmomatic (13), and reads were mapped using BWA against the German PEDV strain L00721 (GenBank LM645057) (14). Remaining gaps were closed using Sanger sequencing.

Isolate 15V010 contained a genome with a size of 28,028 nucleotides (nt), excluding the poly-A tail. Open reading frame 1 (ORF1), encoding the replicase polyprotein, was located between nt 293 and 12601 and between nt 12601 and 20637, with a ribosomal frameshift between both parts. The other genes were arranged as follows: spike (S), nt 20634 to 24785; accessory membrane protein (mp, ORF3), nt 24785 to 25459; envelope (E), nt 25440 and 25670; membrane (M), 25678 to 26358; and nucleocapsid (N), nt 26370 to 27695.

At the whole-genome level, strain 15V010 was most closely related to strain L00721 (99.9% nucleotide similarity) isolated in Germany in 2014, and the prototype U.S. INDEL strain OH851 (99.4%). Strain 15V010 was less closely related to the former prototype European strain CV777 (97.0%).

In conclusion, a PEDV strain from a novel outbreak in Belgium was characterized and showed high relatedness to recent German PEDV isolates and U.S. PEDV INDEL strains. Our findings indicate that PEDV is still or again circulating in Europe. Surveillance and development of prophylactic measures are urgently needed.

Nucleotide sequence accession number. The complete genome sequence of PEDV BEL/15V010/2015 has been deposited at GenBank under the accession number [KR003452](https://www.ncbi.nlm.nih.gov/nuclseq/KR003452).

ACKNOWLEDGMENTS

This work was financially supported by the Laboratory of Virology from Ghent University, and by the Laboratory of Clinical and Epidemiological Virology from the KU Leuven–University of Leuven. S.T. was supported by a fellowship of the Special Research Fund of Ghent University. N.C.N. and I.D.R. were supported by doctoral fellowships from the Institute for the Promotion of Innovation through Science and Technology in Flanders (IWT Vlaanderen).

REFERENCES

1. Martelli P, Lavazza A, Nigrelli AD, Merialdi G, Alborali LG, Pensaert MB. 2008. Epidemic of diarrhoea caused by porcine epidemic diarrhoea virus in Italy. *Vet Rec* 162:307–310. <http://dx.doi.org/10.1136/vr.162.10.307>.
2. Pensaert MB, de Bouck P. 1978. A new coronavirus-like particle associated with diarrhea in swine. *Arch Virol* 58:243–247. <http://dx.doi.org/10.1007/BF01317606>.
3. Luo Y, Zhang J, Deng X, Ye Y, Liao M, Fan H. 2012. Complete genome sequence of a highly prevalent isolate of porcine epidemic diarrhea virus in South China. *J Virol* 86:9551. <http://dx.doi.org/10.1128/JVI.01455-12>.
4. Gao Y, Kou Q, Ge X, Zhou L, Guo X, Yang H. 2013. Phylogenetic analysis of porcine epidemic diarrhea virus field strains prevailing recently in China. *Arch Virol* 158:711–715. <http://dx.doi.org/10.1007/s00705-012-1541-2>.
5. Fan H, Zhang J, Ye Y, Tong T, Xie K, Liao M. 2012. Complete genome sequence of a novel porcine epidemic diarrhea virus in south China. *J Virol* 86:10248–10249. <http://dx.doi.org/10.1128/JVI.01589-12>.
6. Vlasova AN, Marthaler D, Wang Q, Culhane MR, Rossow KD, Rovira A, Collins J, Saif LJ. 2014. Distinct characteristics and complex evolution of PEDV strains, North America, May 2013–February 2014. *Emerg Infect Dis* 20:1620–1628. <http://dx.doi.org/10.3201/eid2010.140491>.
7. Jung K, Wang Q, Scheuer KA, Lu Z, Zhang Y, Saif LJ. 2014. Pathology of US porcine epidemic diarrhea virus strain PC21A in gnotobiotic pigs. *Emerg Infect Dis* 20:662–665. <http://dx.doi.org/10.3201/eid2004.131685>.
8. Marthaler D, Jiang Y, Otterson T, Goyal S, Rossow K, Collins J. 2013. Complete genome sequence of porcine epidemic diarrhea virus strain USA/Colorado/2013 from the United States. *Genome Announc* 1(4): e00555-13. <http://dx.doi.org/10.1128/genomeA.00555-13>.
9. Stevenson GW, Hoang H, Schwartz KJ, Burrough ER, Sun D, Madson D, Cooper VL, Pillatzki A, Gauger P, Schmitt BJ, Koster LG, Killian ML, Yoon KJ. 2013. Emergence of *Porcine epidemic diarrhea virus* in the United States: clinical signs, lesions, and viral genomic sequences. *J Vet Diagn Invest* 25:649–654. <http://dx.doi.org/10.1177/1040638713501675>.
10. Chen Q, Li G, Stasko J, Thomas JT, Stensland WR, Pillatzki AE, Gauger PC, Schwartz KJ, Madson D, Yoon KJ, Stevenson GW, Burrough ER, Harmon KM, Main RG, Zhang J. 2014. Isolation and characterization of porcine epidemic diarrhea viruses associated with the 2013 disease outbreak among swine in the United States. *J Clin Microbiol* 52:234–243. <http://dx.doi.org/10.1128/JCM.02820-13>.
11. Wang L, Byrum B, Zhang Y. 2014. New variant of porcine epidemic diarrhea virus, United States, 2014. *Emerg Infect Dis* 20:917–919. <http://dx.doi.org/10.3201/eid2005.140195>.
12. Hanke D, Jenckel M, Petrov A, Ritzmann M, Stadler J, Akimkin V, Blome S, Pohlmann A, Schirmmeier H, Beer M, Höper D. 2015. Comparison of porcine epidemic diarrhea viruses from Germany and the United States, 2014. *Emerg Infect Dis* 21:493–496. <http://dx.doi.org/10.3201/eid2103.141165>.
13. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <http://dx.doi.org/10.1093/bioinformatics/btu170>.
14. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* 25:1754–1760. <http://dx.doi.org/10.1093/bioinformatics/btp324>.